



The Physics of Electroweak-scale right-handed neutrinos

P. Q. Hung

University of Virginia, Charlottesville

VSOP15, 20-31 July, 2009

Outline

- A brief history of neutrinos.

Outline

- A brief history of neutrinos.
- The neutrino oscillation revolution.

Outline

- A brief history of neutrinos.
- The neutrino oscillation revolution.
- The Standard Model

Outline

- A brief history of neutrinos.
- The neutrino oscillation revolution.
- The Standard Model
- Scenarios of neutrino masses

Outline

- A brief history of neutrinos.
- The neutrino oscillation revolution.
- The Standard Model
- Scenarios of neutrino masses
- Electroweak-scale right-handed neutrinos


Outline

- A brief history of neutrinos.
- The neutrino oscillation revolution.
- The Standard Model
- Scenarios of neutrino masses
- Electroweak-scale right-handed neutrinos
- The LHC as the discovery tool for the origin of masses, including those of neutrinos.

A Brief History of Neutrinos


The Birth of the Neutrino

- The puzzle of β -decays:
The electron (β) energy in the decay
$$A_Z \rightarrow A_{Z+1} + e^-$$
is not equal to the mass difference between A_Z and A_{Z+1} . In fact ...



A Brief History of Neutrinos

- Give up energy conservation?
Wolfgang Pauli's desperate attempt to save it
in 1930:



A Brief History of Neutrinos

- Give up energy conservation?
Wolfgang Pauli's desperate attempt to save it in 1930:
$$A_Z \rightarrow A_{Z+1} + e^- + \textit{unknown particle}$$
- Unknown Particle ("neutron" by Pauli):
electrically neutral, spin 1/2 particle with mass less than 1% of the proton mass.

A Brief History of Neutrinos

- Give up energy conservation?
Wolfgang Pauli's desperate attempt to save it in 1930:
$$A_Z \rightarrow A_{Z+1} + e^- + \textit{unknown particle}$$
- Unknown Particle ("neutron" by Pauli):
electrically neutral, spin 1/2 particle with mass less than 1% of the proton mass.
- In 1932, Chadwick discovered a electrically neutral, spin 1/2 particle with mass comparable to the proton mass: The real neutron and not Pauli's particle.

A Brief History of Neutrinos

- Birth of the Weak Interactions: Enrico Fermi wrote down in 1933 a theory describing β -decay involving the new particle of Pauli and called it the **neutrino** (little neutral one).

A Brief History of Neutrinos

- Birth of the Weak Interactions: Enrico Fermi wrote down in 1933 a theory describing β -decay involving the new particle of Pauli and called it the **neutrino** (little neutral one).
- Neutrinos are **elusive**: light, electrically neutral and interacting weakly with matter. Hard to detect!
If one uses a wall of lead, how thick should it be to stop a beam of neutrinos?

A Brief History of Neutrinos

- Typical low energy (MeV) cross section

$$\sigma \approx 10^{-47} \text{ m}^2.$$

Mean free path for neutrinos going through
e.g. Lead:

- Number density of nucleons in Pb:

$$n = \frac{11400 \text{ kg/m}^3}{1.67 \times 10^{-27} \text{ kg}}.$$

- Number of interaction per meter:

$$\sigma \times n = 10^{-47} \text{ m}^2 \times \frac{11400 \text{ kg/m}^3}{1.67 \times 10^{-27} \text{ kg}}$$

- Mean free path:

$$\lambda = \frac{1}{\sigma \times n} \approx 1.5 \times 10^{17} \text{ m} \approx 1.6 \text{ light years}$$

A Brief History of Neutrinos

- Every second a few **trillions** neutrinos produced from the sun go through our body! And we don't even notice it...

A Brief History of Neutrinos

- Every second a few **trillions** neutrinos produced from the sun go through our body! And we don't even notice it...
- There are about 330 neutrinos $/cm^3$ left over from the Big Bang.

A Brief History of Neutrinos

- Every second a few **trillions** neutrinos produced from the sun go through our body! And we don't even notice it...
- There are about 330 neutrinos $/cm^3$ left over from the Big Bang.
- How are we going to discover **any of that stuff?**

A Brief History of Neutrinos

- Every second a few **trillions** neutrinos produced from the sun go through our body! And we don't even notice it...
- There are about 330 neutrinos $/cm^3$ left over from the Big Bang.
- How are we going to discover **any of that stuff?**
- Huge detector and lots of neutrinos!

A Brief History of Neutrinos

The Age of Discovery

- Fred Reines and Clyde Cowan 1956's experiment: 200 liters of water (lots of protons!), 40 kg of cadmium chloride to capture neutrons, scintillators to detect the photons.

A Brief History of Neutrinos

The Age of Discovery

- Fred Reines and Clyde Cowan 1956's experiment: 200 liters of water (lots of protons!), 40 kg of cadmium chloride to capture neutrons, scintillators to detect the photons.
- Source of neutrinos: Savannah River **nuclear reactor**, South Carolina: ~ 10 trillions neutrinos/ $sec.cm^2$ in the form of $\bar{\nu}_e$.

A Brief History of Neutrinos

- Reaction in detector: $\bar{\nu}_e + p \rightarrow n + e^+$ followed by $e^+ + e^- \rightarrow 2\gamma$ and $n + {}^{108}\text{Cd} \rightarrow {}^{109}\text{Cd}^* \rightarrow {}^{109}\text{Cd} + \gamma$

A Brief History of Neutrinos

- Reaction in detector: $\bar{\nu}_e + p \rightarrow n + e^+$ followed by $e^+ + e^- \rightarrow 2\gamma$ and $n + {}^{108}\text{Cd} \rightarrow {}^{109}\text{Cd}^* \rightarrow {}^{109}\text{Cd} + \gamma$
- Observation: about 3 interactions per hour.
Predicted cross section: $6 \times 10^{-44} \text{ cm}^2$.
Measured cross section: $6.3 \times 10^{-44} \text{ cm}^2$.
Discovery made 26 years after the neutrino was postulated.

A Brief History of Neutrinos

- Reaction in detector: $\bar{\nu}_e + p \rightarrow n + e^+$ followed by $e^+ + e^- \rightarrow 2\gamma$ and $n + {}^{108}\text{Cd} \rightarrow {}^{109}\text{Cd}^* \rightarrow {}^{109}\text{Cd} + \gamma$
- Observation: about 3 interactions per hour.
Predicted cross section: $6 \times 10^{-44} \text{ cm}^2$.
Measured cross section: $6.3 \times 10^{-44} \text{ cm}^2$.
Discovery made 26 years after the neutrino was postulated.
- Nobel Prize in 1995 (Reines). Almost 40 years after the discovery of ν_e !

A Brief History of Neutrinos

- Another neutrino, ν_{μ} , was discovered in an **accelerator** by Leon Lederman, Mel Schwartz and Jack Steinberger in 1962. Nobel Prize 1988, 26 years after its discovery.

A Brief History of Neutrinos

- Another neutrino, ν_{μ} , was discovered in an **accelerator** by Leon Lederman, Mel Schwartz and Jack Steinberger in 1962. Nobel Prize 1988, 26 years after its discovery.
- Discovery of 3rd charged lepton in 1975 by Marty Perl (Nobel Prize 1995). Postulate of a 3rd neutrino ν_{τ} . Observed in 2000 at Fermilab (Nobel Prize in 2020-2030?).

Neutrino Oscillation Revolution

The Age of Astrophysical Neutrinos

- Neutrinos come from outer space too! Lots of them!

Neutrino Oscillation Revolution

The Age of Astrophysical Neutrinos

- Neutrinos come from outer space too! Lots of them!
- ν_e 's produced deep inside the sun carry an energy $\sim 0.1 - 10 \text{ MeV}$.

Neutrino Oscillation Revolution

The Age of Astrophysical Neutrinos

- Neutrinos come from outer space too! Lots of them!
- ν_e 's produced deep inside the sun carry an energy $\sim 0.1 - 10 \text{ MeV}$.
- Small cross section \Rightarrow Big detector.

Neutrino Oscillation Revolution

The Age of Astrophysical Neutrinos

- Neutrinos come from outer space too! Lots of them!
- ν_e 's produced deep inside the sun carry an energy $\sim 0.1 - 10 \text{ MeV}$.
- Small cross section \Rightarrow Big detector.
- Ray Davis's Homestake Experiment in South Dakota: **615 tonne** of tetrachloroethylene (dry cleaning fluid) in a tank 4,800 feet underground.

Neutrino Oscillation Revolution


- 1968 result: neutrino count $\sim 1/3$ of John Bahcall's calculation!
Two possibilities (assuming the experiment is right):
 - The solar model is wrong.
 - Something weird is happening: $2/3$ of ν_e 's disappear on the way to the Earth! (First hint of neutrino mass.)

Neutrino Oscillation Revolution

- 1968 result: neutrino count $\sim 1/3$ of John Bahcall's calculation!
Two possibilities (assuming the experiment is right):
 - The solar model is wrong.
 - Something weird is happening: $2/3$ of ν_e 's disappear on the way to the Earth! (First hint of neutrino mass.)
- This is the so-called **Solar Neutrino Problem**.
What happens to ν_e 's on their way from the **Sun** to **Earth**?


Neutrino Oscillation Revolution

Atmospheric Neutrinos




Neutrino Oscillation Revolution

Super-K Detector, Kamiokande, Japan



Neutrino Oscillation Revolution


- Relativistic muon from e.g. $\nu_{\mu} + n \rightarrow \mu^{-} + p$



Sharp!


Neutrino Oscillation Revolution

- Relativistic muon from e.g. $\nu_{\mu} + n \rightarrow \mu^{-} + p$



Sharp!

- Relativistic electron from e.g. $\nu_e + n \rightarrow e^{-} + p$



Fuzzy! Why?

Neutrino Oscillation Revolution

- Expected $\nu_\mu/\nu_e = 2/1$. Found $\nu_\mu/\nu_e = 1.3/1!$ (1998).

Neutrino Oscillation Revolution

- Expected $\nu_\mu/\nu_e = 2/1$. Found $\nu_\mu/\nu_e = 1.3/1!$ (1998).
- Where did 35 % of ν_μ 's go? Could they change into another neutrino which is not detectable?

Neutrino Oscillation Revolution

- Expected $\nu_\mu/\nu_e = 2/1$. Found $\nu_\mu/\nu_e = 1.3/1!$ (1998).
- Where did 35 % of ν_μ 's go? Could they change into another neutrino which is not detectable?
- How does one describe quantum mechanically the **oscillation** of one quantum state into another?

Neutrino Oscillation Revolution

- Expected $\nu_\mu/\nu_e = 2/1$. Found $\nu_\mu/\nu_e = 1.3/1!$ (1998).
- Where did 35 % of ν_μ 's go? Could they change into another neutrino which is not detectable?
- How does one describe quantum mechanically the **oscillation** of one quantum state into another?
- Definition of **flavor**: $W^+ \rightarrow l_\alpha^+ + \nu_{\alpha}$. l_α^+ : charged lepton mass eigenstate of flavor α (e, μ, τ, \dots)

Neutrino Oscillation Revolution

Simple example of Two Flavor Oscillation

- Flavor eigenstates ν_α ($\alpha = e, \mu$) \neq Mass eigenstates ν_i ($i = 1, 2$).

Neutrino Oscillation Revolution

Simple example of Two Flavor Oscillation

- Flavor eigenstates ν_α ($\alpha = e, \mu$) \neq Mass eigenstates ν_i ($i = 1, 2$).
- $|\nu_e\rangle = \cos\theta|\nu_1\rangle + \sin\theta|\nu_2\rangle$
 $|\nu_\mu\rangle = -\sin\theta|\nu_1\rangle + \cos\theta|\nu_2\rangle$

Neutrino Oscillation Revolution

Simple example of Two Flavor Oscillation

- Flavor eigenstates ν_α ($\alpha = e, \mu$) \neq Mass eigenstates ν_i ($i = 1, 2$).
- $|\nu_e\rangle = \cos\theta|\nu_1\rangle + \sin\theta|\nu_2\rangle$
 $|\nu_\mu\rangle = -\sin\theta|\nu_1\rangle + \cos\theta|\nu_2\rangle$
- At a distance L from the source and for ultrarelativistic neutrinos
 $|\nu_i(L)\rangle = e^{-im_i^2 L/2E} |\nu_i(0)\rangle$

Neutrino Oscillation Revolution

Simple example of Two Flavor Oscillation

- Flavor eigenstates ν_α ($\alpha = e, \mu$) \neq Mass eigenstates ν_i ($i = 1, 2$).
- $|\nu_e\rangle = \cos\theta|\nu_1\rangle + \sin\theta|\nu_2\rangle$
 $|\nu_\mu\rangle = -\sin\theta|\nu_1\rangle + \cos\theta|\nu_2\rangle$
- At a distance L from the source and for ultrarelativistic neutrinos
 $|\nu_i(L)\rangle = e^{-im_i^2 L/2E} |\nu_i(0)\rangle$
- For example $|\nu_e(L)\rangle = \cos\theta e^{-im_1^2 L/2E} |\nu_1\rangle + \sin\theta e^{-im_2^2 L/2E} |\nu_2\rangle$

Neutrino Oscillation Revolution

- Probability for a neutrino of born with flavor α to change into one with one with with a different flavor β at a distance L

$$P(\nu_\alpha \rightarrow \nu_\beta) = \sin^2 2\theta \sin^2 \left(1.267 \frac{\Delta m_{12}^2 L}{E} \frac{\text{GeV}}{\text{eV}^2 \text{ km}} \right)$$

Neutrino Oscillation Revolution

- Probability for a neutrino of born with flavor α to change into one with one with with a different flavor β at a distance L

$$P(\nu_\alpha \rightarrow \nu_\beta) = \sin^2 2\theta \sin^2\left(1.267 \frac{\Delta m_{12}^2 L}{E} \frac{\text{GeV}}{eV^2 \text{ km}}\right)$$

- Probability for it to remain the same or Survival probability:

$$P(\nu_\alpha \rightarrow \nu_\alpha) = 1 - P(\nu_\alpha \rightarrow \nu_\beta)$$

Neutrino Oscillation Revolution

- Probability for a neutrino of born with flavor α to change into one with one with with a different flavor β at a distance L

$$P(\nu_\alpha \rightarrow \nu_\beta) = \sin^2 2\theta \sin^2 \left(1.267 \frac{\Delta m_{12}^2 L}{E} \frac{\text{GeV}}{eV^2 \text{ km}} \right)$$

- Probability for it to remain the same or Survival probability:

$$P(\nu_\alpha \rightarrow \nu_\alpha) = 1 - P(\nu_\alpha \rightarrow \nu_\beta)$$

- Two unknown parameters:

$$\theta \text{ and } \Delta m_{12}^2 = m_1^2 - m_2^2$$

Neutrino Oscillation Revolution

- Probability for a neutrino of born with flavor α to change into one with one with with a different flavor β at a distance L

$$P(\nu_\alpha \rightarrow \nu_\beta) = \sin^2 2\theta \sin^2\left(1.267 \frac{\Delta m_{12}^2 L}{E} \frac{\text{GeV}}{\text{eV}^2 \text{ km}}\right)$$

- Probability for it to remain the same or Survival probability:

$$P(\nu_\alpha \rightarrow \nu_\alpha) = 1 - P(\nu_\alpha \rightarrow \nu_\beta)$$

- Two unknown parameters:

$$\theta \text{ and } \Delta m_{12}^2 = m_1^2 - m_2^2$$

- Choose L and E such that $\Delta m_{12}^2 L/E \approx O(1)$.

Neutrino Oscillation Revolution

- In general, for more than 2 flavors and at $t = 0$

$$|\nu_\alpha(0)\rangle = \sum_i U_{\alpha i}^{\text{lepton}} |\nu_i(0)\rangle$$

Neutrino Oscillation Revolution

- In general, for more than 2 flavors and at $t = 0$
$$|\nu_\alpha(0)\rangle = \sum_i U_{\alpha i}^{\text{lepton}} |\nu_i(0)\rangle$$
- U^{lepton} : Leptonic mixing matrix in charged current interactions similar to the quark Cabibbo-Kobayashi-Maskawa (CKM) matrix.


Neutrino Oscillation Revolution

- In general, for more than 2 flavors and at $t = 0$
 $|\nu_\alpha(0)\rangle = \sum_i U_{\alpha i}^{lepton} |\nu_i(0)\rangle$
- U^{lepton} : Leptonic mixing matrix in charged current interactions similar to the quark Cabibbo-Kobayashi-Maskawa (CKM) matrix.
- At a distance L : $P(\nu_\alpha \rightarrow \nu_\beta) = \sum_{i,j} U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j} \exp\left(-i \frac{\Delta m_{ij}^2 L}{2E}\right)$

Neutrino Oscillation Revolution


- In general, for more than 2 flavors and at $t = 0$
 $|\nu_\alpha(0)\rangle = \sum_i U_{\alpha i}^{lepton} |\nu_i(0)\rangle$
- U^{lepton} : Leptonic mixing matrix in charged current interactions similar to the quark Cabibbo-Kobayashi-Maskawa (CKM) matrix.
- At a distance L : $P(\nu_\alpha \rightarrow \nu_\beta) = \sum_{i,j} U_{\alpha i} U_{\beta i}^* U_{\alpha j}^* U_{\beta j} \exp\left(-i \frac{\Delta m_{ij}^2 L}{2E}\right)$
- To a good approximation, many experiments used the two-flavor oscillation formula. See lectures on neutrinos.

Neutrino Oscillation Revolution




$$P(\bar{\nu}_e \rightarrow \bar{\nu}_e) \text{ in KamLAND}$$

Neutrino Oscillation Revolution




Δm^2 from Atmospheric Neutrinos

Neutrino Oscillation Revolution




Solar neutrinos $\nu_e \rightarrow \nu_X$

Neutrino Oscillation Revolution



Compilation of experimental results

Neutrino Oscillation Revolution



Flavor composition: cross-hatched ν_e ;
left-leaning ν_τ ; right-leaning ν_μ

Neutrino Oscillation Revolution

Summary

- $\Delta m_{atm}^2 \sim 10^{-3} eV^2$; $\Delta m_{solar}^2 \sim 10^{-5} eV^2$

Neutrino Oscillation Revolution

Summary

- $\Delta m_{atm}^2 \sim 10^{-3} eV^2$; $\Delta m_{solar}^2 \sim 10^{-5} eV^2$
- Cosmology: $\sum_i m_i < (0.17 - 2.0)eV$.

Neutrino Oscillation Revolution

Summary

- $\Delta m_{atm}^2 \sim 10^{-3} eV^2$; $\Delta m_{solar}^2 \sim 10^{-5} eV^2$
- Cosmology: $\sum_i m_i < (0.17 - 2.0) eV$.
- Cosmology + oscillation:
 $0.04 eV < m_{heaviest} < (0.07 - 0.7) eV$

Neutrino Oscillation Revolution

Summary

- $\Delta m_{atm}^2 \sim 10^{-3} eV^2$; $\Delta m_{solar}^2 \sim 10^{-5} eV^2$
- Cosmology: $\sum_i m_i < (0.17 - 2.0) eV$.
- Cosmology + oscillation:
 $0.04 eV < m_{heaviest} < (0.07 - 0.7) eV$
- Lepton mixing as deduced from neutrino oscillation data **very different** from quark mixing.

Neutrino Oscillation Revolution

- Lepton mixing:

$$U_{PMNS} \sim \begin{pmatrix} \sqrt{\frac{2}{3}} & \sqrt{\frac{1}{3}} & 0 \\ -\sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & -\sqrt{\frac{1}{2}} \\ -\sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & \sqrt{\frac{1}{2}} \end{pmatrix}$$

Neutrino Oscillation Revolution

- Lepton mixing:

$$U_{PMNS} \sim \begin{pmatrix} \sqrt{\frac{2}{3}} & \sqrt{\frac{1}{3}} & 0 \\ -\sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & -\sqrt{\frac{1}{2}} \\ -\sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & \sqrt{\frac{1}{2}} \end{pmatrix}$$

- Quark mixing with $\lambda \sim 0.2$

$$U_{CKM} \sim \begin{pmatrix} 1 & \lambda & \lambda^3 \\ -\lambda & 1 & \lambda^2 \\ \lambda^3 & -\lambda^2 & 1 \end{pmatrix}$$

Neutrino Oscillation Revolution

The Standard Model

