

# QFT and the EW Standard Model



Just the tip of the iceberg...

Based on lecture notes written with M.A. Vázquez-Mozo


# Apologies

**Never underestimate the pleasure people get  
when they listen to something they already know**

**E. Fermi**

# Outline

- ▶ Why Quantum Field Theory?
- ▶ Quantisation
- ▶ Kinematical symmetries
- ▶ Global symmetries
- ▶ Local symmetries
- ▶ Discrete symmetries
- ▶ Broken symmetries
- ▶ Scale symmetries, renormalisation
- ▶ Standard Model symmetries
- ▶ Amusing examples throughout time permitting



All this in four lectures...

# Do we really need it?

The Schrödinger equation, plus many body physics constructions are very successful in atomic, molecular and solid state physics. The theory of bands, electrical conductivity, atomic bonding, orbitals... are adequately explained in this scheme

$$i \frac{\partial}{\partial t} \Psi(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N, t) = \left( \sum_i \frac{(\mathbf{p}_i - e_i \mathbf{A}_i)}{2m_i} + e_i \Phi_i + V(\mathbf{r}_i) \right) \Psi(\mathbf{r}_j, t)$$
$$P(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N, t) = |\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N, t)|^2, \quad \int \prod_{i=1}^N d^3 \mathbf{r}_i P(\mathbf{r}_i, t) = 1 \quad \forall t$$

---

## A note on conventions

$$\hbar = c = 1, \quad \eta_{\mu\nu} = \text{diag}(+1, -1, -1, -1) \quad \mathbf{F} = \frac{1}{4\pi} \frac{qq'}{r^3} \mathbf{r} \quad \alpha = \frac{e^2}{4\pi\hbar c} \quad e \approx .303$$

# Einstein and Heisenberg complicate our lives

Useful basic formulae. A reminder. Just this once, we reintroduce  $h$  and  $c$

$$p^2 = \left(\frac{E}{c}\right)^2 - m^2 c^2$$

$$E = \pm \sqrt{p^2 c^2 + m^2 c^4} \approx \pm (m c^2 + \frac{p^2}{2m} + \dots)$$

$$\Delta x \Delta p \geq \frac{\hbar}{2}$$

$$\lambda = \frac{h}{mc} \quad \text{Compton wavelength}$$

$$E = \frac{m c^2}{\sqrt{1 - v^2/c^2}} \quad p = \frac{m v}{\sqrt{1 - v^2/c^2}}$$

$$\Delta p \geq mc \quad \Delta E \geq m c^2$$

$$(\Delta x)_{\text{min}} \geq \frac{1}{2} \left( \frac{\hbar}{mc} \right)$$



When the uncertainty in momentum is bigger than  $mc$ , the uncertainty in energy is larger than  $m c^2$ , hence there is enough energy to produce another particle of the same type. In Relativity mass and energy are interchangeable. Hence we cannot localise a particle below its Compton wavelength. If we do, we will not find a single particle, but rather a fairly complicated quantum state with no well-defined number of particles.

Particle production by physical processes should be a central part of the theory.

# Klein paradoxes...



Another way to see the same problem is to consider a particle in a potential barrier in the simplest relativistic generalisation of the Schrödinger equation, the Klein-Gordon equation

$$\left( \frac{\partial^2}{\partial t^2} - \nabla^2 + m^2 \right) \psi(t, \mathbf{x}) = 0$$

$$\begin{aligned} \psi_I(t, x) &= e^{-iEt + ip_1x} + R e^{-iEt - ip_1x}, \\ \psi_{II}(t, x) &= T e^{-iEt + p_2x}, \end{aligned}$$

$$\psi_I(t, 0) = \psi_{II}(t, 0)$$


$$\partial_x \psi_I(t, 0) = \partial_x \psi_{II}(t, 0)$$

$$p_1 = \sqrt{E^2 - m^2},$$

$$p_2 = \sqrt{(E - V_0)^2 - m^2}$$

$$T = \frac{2p_1}{p_1 + p_2},$$

$$R = \frac{p_1 - p_2}{p_1 + p_2}$$




Three cases to consider

- 1)  $E - m > V_0$
- 2)  $E - m < V_0$
- 3)  $V_0 > 2m$
- $V_0 - 2m < E - m < V_0$


In the third case we have the strange situation that we have transmitted wave with negative kinetic energy

$$E - m - V_0$$

In the equation that bears his name, Dirac also found the problem with negative energy states. In his case however he found a rather ingenious way to solve the problem. Since he was describing electrons, he decided to simply fill all the negative energy states, this way Pauli's principle would guarantee stability. His equation also predicted the existence of anti-particles, although at the beginning he was reluctant to accept it. With the Dirac sea we have a simple way to understand anti-electrons = positrons (more later)



## .... Dirac seas



An energetic photon can make a hole. The absence of a negative energy state with negative charge manifests itself as a particle of positive energy and positive charge:

the positron

- ❖ We still have a multi-particle theory after all
- ❖ This does not work for bosons...
- ❖ We should give up the wave equation approach

# Beating a dead horse...


If we still insist against all odds, and decide to violate locality, but to eliminate once and for all the negative energy states by choosing our free Hamiltonian as follows:

$$H = \sqrt{-\nabla^2 + m^2}$$

$$\psi(0, \mathbf{x}) = \delta(\mathbf{x})$$

$$\psi(t, \mathbf{x}) = e^{-it\sqrt{-\nabla^2 + m^2}} \delta(\mathbf{x}) = \int \frac{d^3k}{(2\pi)^3} e^{i\mathbf{k}\cdot\mathbf{x} - it\sqrt{k^2 + m^2}}.$$

$$\psi(t, \mathbf{x}) = \frac{1}{2\pi^2|\mathbf{x}|} \int_{-\infty}^{\infty} k dk e^{ik|\mathbf{x}|} e^{-it\sqrt{k^2 + m^2}}.$$




**Oops!!** we have violated causality! For any  $t > 0$  and any  $|\mathbf{x}|$ , this wave function does not vanish!...



# Relativistic causality

Microscopic causality, Locality in Special Relativity imposes important constraints into what are observables. The light-cone decribes the causal structure of space-time. Physical measurements should be compatible with it



$$[\mathcal{O}(x), \mathcal{O}(y)] = 0, \quad \text{if } (x-y)^2 < 0.$$

- The world is Quantum
- Particle Wave Duality
- Special Relativity
- Microscopic Causality



## LQFT

# From classical to quantum fields

In scattering experiments we observe asymptotic free particles characterised by their energy-momentum charge and other quantum numbers. Consider just E.p. In the NR-case we describe the one-particle states by kets carrying a unitary rep. of the rotation group.

$$|\mathbf{p}\rangle \in \mathcal{H}_1, \quad \langle \mathbf{p} | \mathbf{p}' \rangle = \delta(\mathbf{p} - \mathbf{p}') \quad \mathcal{U}(R)|\mathbf{p}\rangle = |R\mathbf{p}\rangle \quad \hat{p}^i = \int d^3p |\mathbf{p}\rangle p^i \langle \mathbf{p}|$$

To deal with multi-particle states it is convenient to introduce creation and annihilation operators, this leads to the Fock space of states, built out of the vacuum by acting with creation operators:

$$|\mathbf{p}\rangle = a^\dagger(\mathbf{p})|0\rangle, \quad a(\mathbf{p})|0\rangle = 0 \quad \langle 0|0\rangle = 1$$

$$[a(\mathbf{p}), a^\dagger(\mathbf{p}')] = \delta(\mathbf{p} - \mathbf{p}'), \quad [a(\mathbf{p}), a(\mathbf{p}')] = [a^\dagger(\mathbf{p}), a^\dagger(\mathbf{p}')] = 0;$$

We need relativistic invariance, hence we need to find ways to count states in an invariant way. This is necessary also when we deal with decay rates and cross sections. We need to count final states in a way consistent with Lorentz invariance. We can easily construct such an invariant phase space volume:

$$\int \frac{d^4p}{(2\pi)^4} (2\pi) \delta(p^2 - m^2) \theta(p^0) f(p) \quad \text{to integrate over } p_0, \text{ we use a nice identity:}$$

$$\delta[g(x)] = \sum_{x_i = \text{zeros of } g} \frac{1}{|g'(x_i)|} \delta(x - x_i) \quad \delta(p^2 - m^2) = \frac{1}{2p^0} \delta\left(p^0 - \sqrt{\mathbf{p}^2 + m^2}\right) + \frac{1}{2p^0} \delta\left(p^0 + \sqrt{\mathbf{p}^2 + m^2}\right)$$

$$\int \frac{d^3p}{(2\pi)^3} \frac{1}{2E_{\mathbf{p}}} \quad \text{with} \quad E_{\mathbf{p}} \equiv \sqrt{\mathbf{p}^2 + m^2} \quad \text{and} \quad (2E_{\mathbf{p}}) \delta(\mathbf{p} - \mathbf{p}') \quad \text{are invariant}$$

...continued

Now proceed by imitation of the NR case, with the non-trivial result that we have a unitary representation of the Lorentz group

$$|p\rangle = (2\pi)^{\frac{3}{2}} \sqrt{2E_{\mathbf{p}}} |\mathbf{p}\rangle, \quad \langle n|p'\rangle = (2\pi)^3 (2E_{\mathbf{p}}) \delta(\mathbf{p} - \mathbf{p}'), \quad \hat{P}^\mu = \int \frac{d^3p}{(2\pi)^3} \frac{1}{2E_{\mathbf{p}}} |p\rangle p^\mu \langle p|, \quad \mathcal{U}(\Lambda)|p\rangle = |\Lambda^\mu{}_\nu p^\nu\rangle \equiv |\Lambda p\rangle$$

$$\langle 0|0\rangle = 1$$

$$\begin{aligned} \alpha(\mathbf{p}) &\equiv (2\pi)^{\frac{3}{2}} \sqrt{2E_{\mathbf{p}}} a(\mathbf{p}) & [\alpha(\mathbf{p}), \alpha^\dagger(\mathbf{p}')] &= (2\pi)^3 (2E_{\mathbf{p}}) \delta(\mathbf{p} - \mathbf{p}'), & |f\rangle &= \int \frac{d^3p}{(2\pi)^3} \frac{1}{2E_{\mathbf{p}}} f(\mathbf{p}) \alpha^\dagger(\mathbf{p}) |0\rangle \\ \alpha^\dagger(\mathbf{p}) &\equiv (2\pi)^{\frac{3}{2}} \sqrt{2E_{\mathbf{p}}} a^\dagger(\mathbf{p}) & [\alpha(\mathbf{p}), \alpha(\mathbf{p}')] &= [\alpha^\dagger(\mathbf{p}), \alpha^\dagger(\mathbf{p}')] = 0. \end{aligned}$$

Let us construct some observable in this theory. It will be an operator depending on space time, and satisfying some simple conditions:

- ❖ Hermiticity  $\phi(x)^\dagger = \phi(x)$ .
- ❖ Microcausality  $[\phi(x), \phi(y)] = 0, \quad (x - y)^2 < 0$ .
- ❖ Translational invariance  $e^{i\hat{P}\cdot a} \phi(x) e^{-i\hat{P}\cdot a} = \phi(x - a)$
- ❖ Lorentz invariance  $\mathcal{U}(\Lambda)^\dagger \phi(x) \mathcal{U}(\Lambda) = \phi(\Lambda^{-1}x)$ .
- ❖ Linearity  $\phi(x) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{2E_{\mathbf{p}}} [f(\mathbf{p}, x) \alpha(\mathbf{p}) + g(\mathbf{p}, x) \alpha^\dagger(\mathbf{p})]$ .

$$\phi(x) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{2E_{\mathbf{p}}} [e^{-iE_{\mathbf{p}}t + i\mathbf{p}\cdot\mathbf{x}} \alpha(\mathbf{p}) + e^{iE_{\mathbf{p}}t - i\mathbf{p}\cdot\mathbf{x}} \alpha^\dagger(\mathbf{p})]$$

+ve energy
-ve energy

We have obtained from first principles the quantisation of the Klein-Gordon field. There are more straightforward ways, but the procedure shows how to implement the basic principles of the theory, Lorentz invariance, locality and positivity of the spectrum

## Some important properties

$$[\phi(t, \mathbf{x}), \partial_t \phi(t, \mathbf{y})] = i\delta(\mathbf{x} - \mathbf{y}).$$

$$[\phi(x), \phi(x')] = i\Delta(x - x')$$

$$(\partial_\mu \partial^\mu + m^2)\phi(x) = 0$$

$$i\Delta(x - y) = -\text{Im} \int \frac{d^3 p}{(2\pi)^3} \frac{1}{2E_p} e^{-iE_p(t-t') + i\mathbf{p} \cdot (\mathbf{x} - \mathbf{x}'')}$$

$$= \int \frac{d^4 p}{(2\pi)^4} (2\pi) \delta(p^2 - m^2) \varepsilon(p^0) e^{-ip \cdot (x - x')}$$

$$\Delta(x - y) = 0 \quad \text{for } (x - y)^2 < 0$$

The construction is free of paradoxes. It satisfies the KG equation because the +ve and -ve energy plane waves satisfy it. Of course with a free field we do not go very far ...

We should design more powerful techniques leading to similar properties by for more general theories where interactions can take place.

There are two general approaches: the canonical-formalism, and the Feynman path integral. We will briefly introduce the first, just as a reminder.

# Canonical quantisation

Remember: PHYSICS is where the ACTION is!

Proceed by analogy with ordinary QM

$$S[x, \dot{x}] = \int dt L(x, \dot{x})$$
$$L = \sum_i \frac{1}{2} m_i \dot{\mathbf{x}}_i^2 - V(\mathbf{x})$$

$$S[\phi(x)] \equiv \int d^4x \mathcal{L}(\phi, \partial_\mu \phi) = \int d^4x \left( \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{m^2}{2} \phi^2 \right)$$

$$\mathbf{x}_\omega, \dot{\mathbf{x}}_\omega \longleftrightarrow \phi(\mathbf{x}, 0), \dot{\phi}(\mathbf{x}, 0)$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{x}} = \frac{\partial L}{\partial x}$$

$$\partial_\mu \left[ \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi)} \right] - \frac{\partial \mathcal{L}}{\partial \phi} = 0$$

$$\implies (\partial_\mu \partial^\mu + m^2) \phi = 0$$

canonical momenta

$$p = \frac{\partial L}{\partial \dot{x}}$$

$$\pi(x) \equiv \frac{\partial \mathcal{L}}{\partial (\partial_0 \phi)} = \frac{\partial \phi}{\partial t}$$

$$H = \sum_i p_i \dot{x}_i - L$$

$$H \equiv \int d^3x \left( \pi \frac{\partial \phi}{\partial t} - \mathcal{L} \right) = \frac{1}{2} \int d^3x [\pi^2 + (\nabla \phi)^2 + m^2 \phi^2].$$

$$[q^i, p_j] = i\hbar$$

$$[\phi(t, \mathbf{x}), \partial_t \phi(t, \mathbf{y})] = i\delta(\mathbf{x} - \mathbf{y}).$$

Expanding in solutions to the KG equations and performing the canonical quantisation, we recover the algebra of creation and annihilation operator we had before, but we get a surprise

# Casimir effect

Writing the products of creation and ann. operators in NORMAL ORDERING i.e, annihilation operators to the right, we get rid of the sum of the zero point energy of the infinite number of oscillators in the field. In infinite space we subtract it, or simply normal order. When we do not have translational invariance, something interesting happens


$$\hat{H} = \frac{1}{2} \int \frac{d^3 p}{(2\pi)^3} \left[ \hat{\alpha}^\dagger(\mathbf{p}) \hat{\alpha}(\mathbf{p}) + (2\pi)^3 E_{\mathbf{p}} \delta(\mathbf{0}) \right]$$

$$= \int \frac{d^3 p}{(2\pi)^3} \frac{1}{2E_{\mathbf{p}}} E_{\mathbf{p}} \hat{\alpha}^\dagger(\mathbf{p}) \hat{\alpha}(\mathbf{p}) + \frac{1}{2} \int d^3 p E_{\mathbf{p}} \delta(\mathbf{0})$$

$$E(d)_{\text{reg}} = E(d)_{\text{vac}} - E(\infty)_{\text{vac}}$$

The force per unit area is the derivative of this quantity with respect to  $d$  divided by the area of the plates. The result is finite and attractive, the Casimir force! Which has been measured (of course for the electromagnetic field)

$$P_{\text{Casimir}} = -\frac{\pi^2}{240} \frac{1}{d^4}$$



# Lorentz and Poincaré Groups

In trying to systematically construct viable QFTs it is useful to understand the representations of the Lorentz (and Poincaré) groups.

The Hilbert space of states has to carry a unitary representation of the Lorentz group, so that quantum amplitudes are consistent with Unitarity and Relativistic Invariance. The fields themselves however, transform under finite dimensional representations. They are much easier to study. Just recall the usual rotation group SU(2). The Lorentz group, also known as SO(3,1) preserves the Minkowski metric



$$ds^2 = dt^2 - dx^2 - dy^2 - dz^2 = \eta_{\mu\nu} dx^\mu dx^\nu \quad \mu, \nu = 0, 1, 2, 3$$

$$x'^\mu = \Lambda^\mu_\nu x^\nu \quad \eta_{\mu\nu} \Lambda^\mu_\alpha \Lambda^\nu_\beta = \eta_{\alpha\beta}$$

$$\det \Lambda = \pm 1 \quad (\Lambda^0_0)^2 - \sum_{i=1}^3 (\Lambda^i_0)^2 = 1$$

- $\mathcal{L}^\uparrow_+$ : proper, orthochronous transformations with  $\det \Lambda = 1, \Lambda^0_0 \geq 1$ .
- $\mathcal{L}^\uparrow_-$ : improper, orthochronous transformations with  $\det \Lambda = -1, \Lambda^0_0 \geq 1$ .
- $\mathcal{L}^\downarrow_-$ : improper, non-orthochronous transformations with  $\det \Lambda = -1, \Lambda^0_0 \leq -1$ .
- $\mathcal{L}^\downarrow_+$ : proper, non-orthochronous transformations with  $\det \Lambda = 1, \Lambda^0_0 \leq -1$ .

$$\mathcal{L}^\uparrow_+ \xrightarrow{\mathcal{P}} \mathcal{L}^\uparrow_-, \quad \mathcal{L}^\uparrow_+ \xrightarrow{\mathcal{T}} \mathcal{L}^\downarrow_-, \quad \mathcal{L}^\uparrow_+ \xrightarrow{\mathcal{PT}} \mathcal{L}^\downarrow_+$$

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \quad \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

# Lorentz and Poincaré Groups

$$R(\mathbf{e}, \varphi) = e^{-i\varphi \mathbf{e} \cdot \mathbf{J}}$$

$$B(\mathbf{u}, \lambda) = e^{-i\lambda \mathbf{u} \cdot \mathbf{M}}$$

Rotations and boosts generate Lorentz transformation, hence six parameter and six generators of infinitesimal transformations. The algebra is easy to obtain and “diagonalise”

$$[J_i, J_j] = ie_{ijk} J_k,$$

$$[J_i, M_k] = ie_{ijk} M_k,$$

$$[M_i, M_j] = -ie_{ijk} J_k$$

$$J_k^\pm = \frac{1}{2}(J_k \pm iM_k)$$

$$[J_i^\pm, J_j^\pm] = ie_{ijk} J_k^\pm,$$

$$[J_i^+, J_j^-] = 0.$$

The representations of each SU(2) are labelled by a single integer or half integer “angular” momentum  $s=0, 1/2, 1, 3/2, \dots$ . Under parity

$$(s_+, s_-)$$

Representation	Type of field
$(0, 0)$	Scalar
$(\frac{1}{2}, 0)$	Right-handed spinor
$(0, \frac{1}{2})$	Left-handed spinor
$(\frac{1}{2}, \frac{1}{2})$	Vector
$(1, 0)$	Selfdual antisymmetric 2-tensor
$(0, 1)$	Anti-selfdual antisymmetric 2-tensor

$$\begin{array}{ccc} \mathbf{J} & \xrightarrow{P} & \mathbf{J} \\ \mathbf{M} & \rightarrow & -\mathbf{M} \\ \mathbf{J}^\pm & \rightarrow & \mathbf{J}^\mp \\ (s_+, s_-) & \rightarrow & (s_-, s_+) \end{array}$$

$$\mathbf{J} = \mathbf{J}^+ + \mathbf{J}^-$$

$$(s_+, s_-) = \sum_{j=|s_+|}^{s_++s_-} \mathbf{j}$$



# Weyl spinors

The simplest representations have fundamental physical importance, they are called Weyl spinors. Clearly they are representations of the connected component of  $SO(3,1)$ , but not of parity, since parity interchanges the representations

$$J_i^+ = \frac{1}{2}\sigma_i,$$


$$J_i^- = 0$$

for  $(\frac{1}{2}, \mathbf{0})$ ,

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$\sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$

$$\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$



$$J_i^+ = 0,$$

$$J_i^- = \frac{1}{2}\sigma_i$$

for  $(\mathbf{0}, \frac{1}{2})$ .

$$u_{\pm} \longrightarrow e^{-\frac{i}{2}(\theta \mathbf{n} \mp i\beta) \cdot \sigma} u_{\pm}$$

$$u_{\pm} \longrightarrow e^{i\theta} u_{\pm}$$

Consider for simplicity this global symmetry: fermion number

$$\sigma_{\pm}^{\mu} = (\mathbf{1}, \pm\sigma_i)$$

$$u_{+}^{\dagger} \sigma_{+}^{\mu} u_{+}$$

$$\mathcal{L}_{\text{Weyl}}^{\pm} = i u_{\pm}^{\dagger} (\partial_t \pm \sigma \cdot \nabla) u_{\pm} = i u_{\pm}^{\dagger} \sigma_{\pm}^{\mu} \partial_{\mu} u_{\pm}$$

$$(\partial_0 \pm \sigma \cdot \nabla) u_{\pm} = 0$$


$$u_{\pm}(x) = u_{\pm}(k) e^{-ik \cdot x}$$

$$(|\mathbf{k}| \mp \mathbf{k} \cdot \sigma) u_{\pm} = 0$$

$$k_0^2 = k_0^2 - \mathbf{k}^2 = 0$$

$$u_{+} : \quad \frac{\sigma \cdot \mathbf{k}}{|\mathbf{k}|} = 1,$$

$$u_{-} : \quad \frac{\sigma \cdot \mathbf{k}}{|\mathbf{k}|} = -1$$



positive helicity, right handed antineutrinos

negative helicity, left handed, neutrinos

# Charge conjugation and Majorana masses

We know that under parity, the L,R Weyl spinors are exchanged. Another way to exchange them is via complex conjugation, later to be related to charge conjugation

$$\begin{aligned}
 M_L &= e^{-\frac{i}{2}\theta \cdot \sigma - \frac{1}{2}\beta \cdot \sigma} & \det M_L &= 1 & \det M &= \epsilon_{ab} M_{a1} M_{b2} & \epsilon &= i\sigma_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \\
 M_R &= e^{-\frac{i}{2}\theta \cdot \sigma + \frac{1}{2}\beta \cdot \sigma} & \det M_R &= 1 & \det M \epsilon_{ab} &= \epsilon_{cd} M_{ca} M_{db}
 \end{aligned}$$

Using  $\sigma^* = -\sigma_2 \sigma \sigma_2$

$$\begin{aligned}
 \psi_L^C &= \sigma_2 \psi_L^* & \text{transforms like } \psi_R \\
 \psi_R^C &= \sigma_2 \psi_R^* & \text{transforms like } \psi_L
 \end{aligned}$$

► We can express any theory fully in terms of L or R fermions.

$$\begin{aligned}
 \mathcal{L}_{\text{Weyl}}^\pm &= i u_\pm^\dagger \sigma_\pm^\mu \partial_\mu u_\pm + \frac{m}{2} \left( \epsilon_{ab} u_\pm^a u_\pm^b + \text{h.c.} \right) \\
 \epsilon_{ab} u^a u^b &= u^1 u^2 - u^2 u^1
 \end{aligned}$$

► Charge conjugation and parity exchange L and R

Most general Majorana mass, Takagi factorisation

► A parity invariant theory requires L,R spinors at the same time

$$\frac{1}{2} (M_{IJ} \epsilon_{ab} u^{a,I} u^{b,J} + \text{h.c.}),$$

$I, J = 1, \dots, N_F$ ,  $M_{IJ} = M_{JI}$  complex

► We can construct a mass for pure L spinors if we ignore fermion number

$$M = U \begin{pmatrix} m_1 & \dots & 0 \\ 0 & \ddots & 0 \\ 0 & \dots & m_{N_F} \end{pmatrix} U^T$$

► Fermions are anticommuting  $m_i$  are positive square roots of  $MM^\dagger$

This is the most general fermion mass matrix!!! It includes CKM, in fact it is more general

# Weyl + parity: Dirac

$$\left(\frac{1}{2}, \mathbf{0}\right) \oplus \left(\mathbf{0}, \frac{1}{2}\right)$$

$$P: u_{\pm} \longrightarrow u_{\mp} \quad \psi = \begin{pmatrix} u_+ \\ u_- \end{pmatrix} \quad \left. \begin{array}{l} i\sigma_+^{\mu} \partial_{\mu} u_+ = m u_- \\ i\sigma_-^{\mu} \partial_{\mu} u_- = m u_+ \end{array} \right\} \implies i \begin{pmatrix} \sigma_+^{\mu} & 0 \\ 0 & \sigma_-^{\mu} \end{pmatrix} \partial_{\mu} \psi = m \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \psi$$

$$\gamma^{\mu} = \begin{pmatrix} 0 & \sigma_-^{\mu} \\ \sigma_+^{\mu} & 0 \end{pmatrix} \quad \bar{\psi} \equiv \psi^{\dagger} \gamma^0 = \psi^{\dagger} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \mathcal{L}_{\text{Dirac}} = \bar{\psi} (i\gamma^{\mu} \partial_{\mu} - m) \psi$$

## DIRAC LOGY

$$\{\gamma^{\mu}, \gamma^{\nu}\} = 2\eta^{\mu\nu} \quad \gamma_5 = -i\gamma^0\gamma^1\gamma^2\gamma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad P_{\pm} = \frac{1}{2}(1 \pm \gamma_5) \quad \begin{array}{l} P_+ \psi = \begin{pmatrix} u_+ \\ 0 \end{pmatrix} \\ P_- \psi = \begin{pmatrix} 0 \\ u_- \end{pmatrix} \end{array}$$

$$\text{Tr } \gamma^{\mu} \gamma^{\nu} = 4\eta^{\mu\nu}$$

$$\text{Tr } \gamma^{\mu} \gamma^{\nu} \gamma^{\alpha} \gamma^{\beta} = 4\eta^{\mu\nu} \eta^{\alpha\beta} - 4\eta^{\mu\alpha} \eta^{\beta\nu} + 4\eta^{\mu\beta} \eta^{\alpha\nu}$$

$$\text{Tr } \gamma_5 \gamma^{\alpha} \gamma^{\beta} \gamma^{\mu} \gamma^{\nu} = 4i \epsilon^{\alpha\beta\mu\nu}$$

$$u(\mathbf{k}, s) e^{-ik \cdot x},$$

$$(\not{k} - m)u(\mathbf{k}, s) = 0.$$

$$v(\mathbf{k}, s) e^{ik \cdot x}$$

$$(\not{k} + m)v(\mathbf{k}, s) = 0$$

$$k^2 = m^2$$

We look for +ve and -ve energy solutions as usual

$$\begin{array}{ll} \bar{u}(\mathbf{k}, s) u(\mathbf{k}, s) = 2m, & \bar{v}(\mathbf{k}, s) v(\mathbf{k}, s) = -2m, \\ \bar{u}(\mathbf{k}, s) \gamma^{\mu} u(\mathbf{k}, s) = 2k^{\mu}, & \bar{v}(\mathbf{k}, s) \gamma^{\mu} v(\mathbf{k}, s) = 2k^{\mu}, \\ \sum_{s=\pm\frac{1}{2}} u_{\alpha}(\mathbf{k}, s) \bar{u}_{\beta}(\mathbf{k}, s) = (\not{k} + m)_{\alpha\beta} & \sum_{s=\pm\frac{1}{2}} v_{\alpha}(\mathbf{k}, s) \bar{v}_{\beta}(\mathbf{k}, s) = (\not{k} - m)_{\alpha\beta} \end{array}$$